

Extracorporeal Treatment in Phenytoin Poisoning: Systematic Review and Recommendations from the EXTRIP Workgroup

Kurt Anseeuw, MD, MSc,¹ James B. Mowry, PharmD,² Emmanuel A. Burdmann, MD PhD,³ Marc Ghannoum, MDCM,⁴ Robert S. Hoffman, MD,⁵ Sophie Gosselin, MD,⁶ Valery Lavergne, MD, MSc,⁷ and Thomas D. Nolin, PharmD, PhD,^{8,*} on behalf of the EXTRIP Workgroup

Author affiliations:

¹ Campus Stuivenberg, Emergency Medicine, Antwerpen, Belgium; ² Indiana University Health, Indiana Poison Center, Indianapolis, IN, USA; ³ LIM 12, University of Sao Paulo Medical School, Division of Nephrology, Sao Paulo, Brazil; ⁴ Department of Nephrology, Verdun Hospital, University of Montreal, Verdun, QC, Canada; ⁵ Division of Medical Toxicology, Ronald O. Perelman Department of Emergency Medicine, New York University School of Medicine, New York, NY, USA; ⁶ Department of Emergency Medicine, Medical Toxicology Division, McGill University Health Centre & Department of Medicine McGill University, Montreal, QC, Canada; ⁷ Department of Medical Biology, Sacre-Coeur Hospital, University of Montreal, Montreal, QC, Canada; ⁸ Department of Pharmacy and Therapeutics, and Department of Medicine Renal Electrolyte Division, University of Pittsburgh Schools of Pharmacy and Medicine, Pittsburgh, PA, USA

***Corresponding author:** Thomas D. Nolin, PharmD, PhD
University of Pittsburgh School of Pharmacy
Department of Pharmacy and Therapeutics
208 Salk Pavilion, 335 Sutherland Drive
Pittsburgh, PA 15260 USA
TEL: 1.412.624.4683
E-mail: nolin@pitt.edu

Article Category: Special Report

Short Title: Extracorporeal Treatment in Phenytoin Poisoning

Word Count: Abstract: 260 Text: 4,048

References: 115 **# Figures:** 1 **# Tables:** 5 **#Box :** 1 **# Suppl Items:** 1

Index Words: Phenytoin; poisoning; toxicity; EXTRIP (Extracorporeal Treatments in Poisoning); recommendations; indications; extracorporeal treatment (ECTR); hemodialysis; renal replacement therapy; pharmacokinetics; toxicokinetics; dialyzability

Support and Financial Disclosure Declaration

Funding: Funding for EXTRIP was obtained from industry in the form of unrestricted educational grants. These funds were used solely for expenses related to literature retrieval, translation of publications, and for reimbursement of conference calls and travel expenses for attendance at EXTRIP meetings. Current EXTRIP sponsors are Leo Pharma (\$40,000), Janssen-Ortho (\$30,000), Fresenius Canada (\$20,000), Amgen Canada (\$40,000), and Servier (\$3,000). There was no industry input into scientific content, development, or publication of the recommendations. Furthermore, industry presence at meetings was not allowed, nor was industry awareness or comment on the recommendations accepted. A list of EXTRIP sponsors can be found at www.extrip-workgroup.org.

Financial Disclosure: Competing interest from participants include: SKA: Advisory board (Merck), MG: Lecturer (Amgen Canada and Janssen-Ortho), advisory boards (Genzyme, Amgen Canada), RM: Consultative work (Baxter Healthcare), Travel costs unrelated to EXTRIP (Roche, Amgen), JK: Advisory board and grant (Fresenius Medical Care), DG: Consultant (Takeda), Honorarium (Genzyme, Mission Pharmacal), Advisory board (Retrophin, Astra Zeneca), Owner (The Ravine Group), Research site collaborator (Reata, Amgen, Hospira), RM: Grant (Hospira), TB: Guest speaker (Gambro), KL: Stock (Amgen). The remaining authors declare that they have no competing interests. Complete financial disclosure for each EXTRIP member can be found at www.extrip-workgroup.org.

ABSTRACT

The Extracorporeal Treatments in Poisoning (EXTRIP) Workgroup conducted a systematic literature review using a standardized process to develop evidence-based recommendations on the use of extracorporeal treatment (ECTR) in patients with phenytoin poisoning. The authors reviewed all articles, extracted data, summarized findings, and proposed structured voting statements following a pre-determined format. A two-round modified Delphi method was used to reach a consensus on voting statements, and the RAND/UCLA Appropriateness Method was employed to quantify disagreement. 51 articles met inclusion criteria. Only case reports, case series, and pharmacokinetic studies were identified yielding a very low quality of evidence. Clinical data from 31 patients and toxicokinetic grading from 46 patients were abstracted. The Workgroup concluded that phenytoin is moderately dialyzable (Level of evidence = C) despite its high protein binding and made the following recommendations: ECTR would be reasonable in selected cases of severe phenytoin poisoning, (Neutral recommendation, 3D). ECTR is suggested if prolonged coma is present or expected (2D) and it would be reasonable if prolonged incapacitating ataxia is present or expected (3D). If ECTR is used, it should be discontinued when clinical improvement is apparent (1D). The preferred ECTR modality in phenytoin poisoning is intermittent hemodialysis (1D), but hemoperfusion is an acceptable alternative if hemodialysis is not available (1D). In summary, phenytoin appears to be amenable to extracorporeal removal. However, because of the low incidence of irreversible tissue injury or fatality related to phenytoin poisoning, and the relatively limited effect of ECTR on phenytoin removal, the Workgroup proposed the use of ECTR only in very select patients with severe phenytoin poisoning.

INTRODUCTION

The Extracorporeal Treatments In Poisoning (EXTRIP) workgroup is composed of international experts representing diverse specialties and professional societies (Item S1) to provide recommendations on the use of extracorporeal treatments (ECTRs) in poisoning (www.extrip-workgroup.org). The rationale, background, objectives, methodology, and its initial recommendations have been published previously.¹⁻¹³ We present a systematic literature review and evidence-based recommendations for the use of ECTR in phenytoin poisoning.

Pharmacology and Toxicokinetics

Phenytoin is a hydantoin derivative that is used as a first-line agent in the control of tonic-clonic and psychomotor seizures, and the prevention and treatment of seizures associated with neurosurgery.¹⁴⁻¹⁶ The main site of action of phenytoin is the motor cortex, where it stabilizes transmembrane flux of ions and reduces post-tetanic potentiation of synapses.¹⁴ Specifically, phenytoin inhibits sodium channels by reducing their capacity for recovery after inactivation.^{14,17} Phenytoin also increases brain concentrations of gamma-aminobutyric acid (GABA), which has an inhibitory action in the cerebral cortex.^{14,15}

Phenytoin has a molecular mass of 252 Da, and it binds extensively to plasma proteins (binding = 90%), a percentage that stays unchanged after overdose^{18,19} but decreases slightly to 75-80% in patients with kidney failure, hypoalbuminemia or cytochrome P450 (CYP) 2C9 genetic polymorphism.²⁰ The unbound or "free" form is responsible for its clinical and toxicological effects.^{21,22} The reported time to peak plasma concentrations in therapeutic dosing is 1.5 to 3 hours for standard formulations and 4 to 12 hours for extended-release formulations. However, oral absorption of phenytoin is slow and variable, and can be delayed and unpredictable during

overdose. Peak plasma concentrations have been observed up to 96 hours after ingestion in the overdose setting.²³⁻²⁵

Phenytoin had a volume of distribution of 0.6-0.8 L/kg, and is predominantly metabolized by the CYP enzyme system to inactive metabolites. The drug exhibits Michaelis-Menten kinetics; as such, increased doses may produce a larger than expected increase in plasma concentrations and prolonged elimination.^{14,21,26} Less than 1% of phenytoin is eliminated unchanged in urine, although its metabolites, including 5-(p-hydroxyphenyl)-5-phenylhydantoin (HPPH), are renally excreted. At therapeutic concentrations, phenytoin's endogenous clearance is 23 mL/min²⁷ and its apparent elimination half-life is approximately 22 hours (range 7 to 42 hours).^{14,21} In overdose, the elimination half-life increases; in one case it was reported to be as long as 103 hours.²⁸ This explains why massive phenytoin ingestions may lead to prolonged toxicity and extended hospital stays. The physicochemical characteristics and pharmacokinetic properties of phenytoin are presented in Table 1.

Overview of Phenytoin Poisoning

US Poison Control Centers documented 2,850 phenytoin exposures in 2013, 528 of which had a clinical outcome defined as moderate outcome or worse, including one fatality.²⁹ Oral overdose is characterized by cerebellar and vestibular effects including multi-directional nystagmus, dizziness, nausea, vomiting and ataxia.^{30,31} Severe overdose may result in coma and marked respiratory depression.^{30,31} To our knowledge, there is no previously published literature on the frequency of clinical effects for phenytoin overdoses. In response, we performed a search in the National Poison Data System from 2000 to 2014 for single substance phenytoin exposures, coded with a serious outcome (major effects or fatalities).³² Of the 734 retrieved cases, respiratory arrest

was reported in 3.1%, respiratory depression in 5.7%, coma in 16.1% and ataxia in 25.1%. The other most common signs and symptoms include seizures (44.1%), drowsiness/lethargy (39%), confusion (23.2%), nystagmus (17.8%), agitation/irritable (15.4%), hypotension (12.5%) and slurred speech (11.4%). Cardiac arrest was present in 3.5% of the cases and 29 patients died.

Mortality or irreversible injury following phenytoin poisoning is infrequent but still reported.³³⁻³⁵ Intravenous overdose produces similar systemic effects to oral overdose, but cardiotoxicity, including hypotension, bradycardia, dysrhythmias and even asystole can occur.^{36,37} These side effects are thought to be caused by the diluent (propylene glycol) rather than phenytoin itself.^{38,39}

Fosphenytoin, the intravenous prodrug of phenytoin, is designed with an extra phosphate linkage enhancing its water solubility. It comes as an injection, dissolved in sterile water and tromethamine buffer. In vivo, fosphenytoin is converted into phenytoin by losing the phosphate group. Though there is limited information on the toxicity of fosphenytoin, data from US poison centers from 2000 to 2014 on 208 single substance fosphenytoin exposures showed the most common symptoms including seizures (22.6%), drowsiness/lethargy (22.6%), no symptoms (17.3%), hypotension (14.9%), other (13.0%), ataxia (11.1%), agitated/irritable (9.1%), confusion (8.2%), vomiting (8.2%) and nystagmus (7.7%).³² In the 25 cases classified as serious (major effects or death), seizures remained the most common symptom at 44.0% followed by hypotension (32.0%), bradycardia (24%), and respiratory arrest (20%). Cardiac arrest was noted in 12.0% of serious cases. Case reports demonstrate that the most serious, acute complication of massive IV fosphenytoin overdoses is cardiovascular in nature (hypotension, bradydysrhythmias, conduction disturbances and even asystole). The signs and symptoms of a classic phenytoin poisoning (coma, ataxia, drowsiness, seizures) manifest as the toxicity progresses.⁴⁰⁻⁴⁵

The onset of symptoms usually occurs within minutes of intravenous administration and within 1 to 2 hours of ingestion, although the latter can be delayed or prolonged for up to a week due to prolonged absorption and saturable metabolism. Factors on admission that correlate with length of stay include disorientation and unarousability, concurrent phenothiazine usage and liver disease,⁴⁶ while morbidity is correlated to ataxia.⁴⁷ While therapeutic concentrations of total plasma phenytoin are reported to be 10-20 mcg/mL, free phenytoin concentrations (1-2 mcg/mL therapeutic; > 5 mcg/mL toxic) are more accurate in predicting clinical effects.⁴⁸ However, there is little correlation between the phenytoin concentration at presentation and the severity of toxicity or length of stay.⁴⁷

The management of patients with phenytoin toxicity is largely supportive, including airway protection and correction of hypotension with intravenous fluids. Advanced cardiac life support guidelines should be followed when indicated. Gastrointestinal decontamination (e.g., single dose activated charcoal) should be given if the patient presents shortly after ingestion and has no contraindications, although there is no evidence that this alters the clinical course.⁴⁹ There are no antidotes available to reverse phenytoin's effects. While multiple-dose activated charcoal (MDAC) increases elimination of phenytoin,⁵⁰⁻⁵² the data are conflicting regarding the improvement of clinical outcome in phenytoin overdose. In some studies MDAC failed to demonstrate a beneficial effect on time to resolution of phenytoin toxicity,⁵³⁻⁵⁵ while in other studies MDAC seems to improve the clinical course.⁵⁶ For the moment, MDAC is not routinely recommended for the treatment of phenytoin ingestions,⁵⁷ but may be considered in selected cases.^{58,59} Current toxicology resources do not routinely recommend ECTR for phenytoin poisoning, claiming unproven benefit on clinical outcomes.⁶⁰⁻⁶²

METHODS

A Predetermined methodology, incorporating recommendations from The Appraisal of Guidelines for Research and Evaluation (AGREE)⁶³ and Grading of Recommendation Assessment, Development and Evaluation (GRADE),⁶⁴ was used and is described in detail elsewhere.² The primary literature search was conducted on July 12th 2012 in Medline, Embase and Cochrane library (Review and Central).

The search strategy was: [phenytoin OR dilantin] AND [toxicity OR poison* OR intoxication OR overdos*] AND [hemoperfusion OR haemoperfusion OR hemofiltration OR haemofiltration OR hemodialysis OR haemodialysis OR hemodiafiltration OR haemodiafiltration OR dialysis OR plasmapheresis OR plasma exchange OR exchange transfusion OR CRRT OR renal replacement therapy].

A manual search of conference proceedings of the European Association of European Association of Poisons Centres and Clinical Toxicologists (EAPCCT) and the North American Congress of Clinical Toxicology (NACCT) annual meetings (2002-2014), and Google Scholar was performed, as well as the bibliography of each article obtained during the literature search.

A subgroup of EXTRIP completed the literature search, reviewed each article, extracted data and summarized findings. The level of evidence assigned to each clinical recommendation (Table 2) and dialyzability were determined based on established criteria.² The potential benefits of the procedure were weighed against its cost, availability, related complications, and alternative treatments. All this information was submitted to the entire workgroup for consideration, along with structured voting statements based on a pre-determined format.

The strength of recommendations was evaluated by a two-round modified Delphi method for each proposed voting statement and a RAND/UCLA Appropriateness Method was used to quantify disagreement between voters, as previously described.^{2,65} Anonymous votes with comments were sent to the epidemiologist who then compiled and returned a summary to each participant. The workgroup met in person to exchange ideas and debate statements. A second vote was later conducted and these results were used in determining the core EXTRIP recommendations. The literature search was updated on November 15th 2014 using the methodology described above; new articles and the updated data summary were submitted to every participant who then updated their votes.

RESULTS

Results of the literature search are presented in Figure 1. A total of 546 articles were identified after removal of duplicates. In the final analysis, 51 studies were included for qualitative analysis: 30 case-reports or case-series (31 patients),^{18,19,33,35-37,66-89} 17 pharmacokinetic studies (54 patients),⁹⁰⁻¹⁰⁶ 1 animal experiment,¹⁰⁷ and 3 *in vitro* studies.¹⁰⁸⁻¹¹⁰ No randomized-controlled trial (RCT) or observational studies were identified.

Clinical Outcomes

The evidence of a clinical effect of ECTR in phenytoin poisoning is comprised only of case reports and case series, which are inherently anecdotal, limited by a lack of controls, and susceptible to publication bias. Therefore, the quality of the evidence for all recommendations was graded as **very low**.

Clinical data from 30 reports and 31 patients were retrieved; the first reported case of ECTR was published in 1958 when hemodialysis was used to treat a boy poisoned with phenytoin.⁶⁶ An aggregate description of clinical outcomes of reported cases is presented in Table 3. The average phenytoin ingestion and peak total concentration were 6.8 grams and 69.8 mcg/mL, respectively. The majority of patients presented with some level of impaired consciousness, and most reported cases were treated using either hemodialysis or hemoperfusion.

In the studied cohort, most patients experienced some improvement during or shortly after ECTR, which was occasionally dramatic when using an efficient ECTR.^{19,71,77,79,87} Conversely, some patients experienced no apparent benefit from ECTR and developed a chronic protracted course or long-term sequelae.^{33,35,66,80,84} In others, incapacitating ataxia was still present one week after exposure.^{70,81} Although the natural history of severely poisoned patients not treated with ECTR suggests survival,^{111,112} irreversible neurological conditions may incur in those most at risk.²³⁻²⁵ In our cohort, survival was noted in some patients who reportedly ingested 10 grams or greater,^{33,67,69,71,84} and who had peak total phenytoin concentrations over 90 mcg/mL.^{33,35,66,70,75,87,88} Complications associated with ECTR included thrombocytopenia after hemoperfusion,^{18,88} and peritonitis following peritoneal dialysis that led to death.⁷⁴ The other fatality appeared unrelated to phenytoin toxicity and followed bowel infarction.⁸⁶

Dialyzability

Phenytoin is a small molecule (252 Da) and has a small volume of distribution of 0.6 to 0.8 L/kg.^{18,19,27} However, because of its extensive binding to plasma proteins (90%), hemoperfusion (HP) and therapeutic plasma exchange (TPE) would theoretically be most likely to efficiently remove phenytoin. TPE can readily remove phenytoin from the vascular compartment, albeit at a

slow rate; on average TPE removes 5-10% of total body load of phenytoin during a single 2-3 hour exchange^{93,95,99,102} and can provide clearances up to 20 mL/min (Table 4).^{78,113} Initial clearances with charcoal HP surpass this range but are subsequently limited by saturation of the column, which usually occurs within 2 hours.^{72,114} Although some sorbent adsorption columns have been tested *in vitro*, their application in clinical practice is unknown.¹⁰⁹

Historical reports suggest a lack of an effect of diffusive techniques including intermittent hemodialysis (HD) because of phenytoin's significant protein binding.^{36,67,71,76,90,92,94} However, encouraging results have been shown with high-efficiency filters,¹¹⁰ especially in patients presenting with conditions known to reduce protein binding (e.g., hypoalbuminemia, kidney disease); clearance may exceed 100 mL/min in uremic patients.¹⁰³ Clearances are inferior but nevertheless considerable in patients who have normal protein binding, although the workgroup acknowledged that additional study is required to confirm this.^{19,86} One report describes a patient who ingested 3.6 grams of phenytoin who was treated with HD.¹⁹ A total of 547 mg of phenytoin was extracted in approximately 6 hours. Overall, the aggregate ECTR clearance (Table 4) and the toxicokinetic grading of all patients (Table 5) demonstrate the superiority of both HP and high-efficiency HD over all other techniques. The advantages of HP over HD for phenytoin removal are less clear; in patients who underwent both techniques, HP was superior to HD in one report,⁸² and inferior in the other.⁸⁶ The addition of a charcoal column in series after a dialysis filter appears to enhance clearance of both HP or HD alone, but requires further study.⁸⁶

Supplementation of albumin into the dialysate does not appear to substantially enhance removal of phenytoin.^{74,110} The data on other liver support therapies, like molecular adsorbent recirculating systems (MARS), are limited: in one case, the apparent phenytoin elimination half-life during MARS was 8.4 h, which is not shorter than what can be achieved with more conventional

and less expensive alternatives.³⁷ Further studies are needed to confirm the role of liver support therapies, especially considering their cost. Other ECTRs show limited phenytoin clearance (< 10 mL/min)¹¹³ including exchange transfusion (ET),⁷⁶ peritoneal dialysis (PD),^{68,69} and continuous renal replacement therapy (CRRT)^{100,104} and are therefore of very limited use in phenytoin poisoning.

In summary, the best reported clearances that can be sustained are with HD, possibly in-series with a charcoal cartridge. Based on the criteria established previously,⁹ the dialyzability of phenytoin is "slightly dialyzable" or "not dialyzable" for less efficient techniques like ET, PD, CRRT, and conventional HD techniques using less efficient cuprophane membranes. Phenytoin is "slightly" to "moderately dialyzable" with TPE and "moderately dialyzable" to "dialyzable" for HP, high-efficiency HD, and liver support therapies, especially if the patient presents with a condition that is associated with decreased protein binding (e.g., hypoalbuminemia, malnutrition, kidney disease). The workgroup preferred a conservative grading and therefore agreed with the following statement: ***Phenytoin is moderately dialyzable (level of evidence = C).*** Again, it was acknowledged that more studies using contemporary ECTR technology and parameters as well as performing more complete toxicokinetic measurements, especially quantification of removal in effluent fluid,⁹ are needed to support this observation.

There are no studies that specifically compare ECTR to MDAC with respect to enhanced elimination of phenytoin. In three case reports, ECTR appeared superior, although the toxicokinetic data are relatively incomplete and cannot be reliably interpreted.^{19,33,84}

RECOMMENDATIONS

An executive summary of the recommendations is provided in **Box 1**.

1. General statement regarding use of ECTR

- ECTR would be reasonable in selected cases of severe phenytoin poisoning (Neutral recommendation - 3D)

Rationale: Phenytoin is a widely used pharmaceutical, and toxicity following an acute ingestion or in therapeutic dosing is common.²⁹ Life-threatening symptoms are infrequent and usually resolve completely with appropriate supportive treatment. No antidotes currently exist to reverse the toxic effects of phenytoin and the use of MDAC remains controversial.^{50,55} In severe cases, incapacitating and prolonged ataxia may occur, which can progress to stupor and coma. ECTR not only removes phenytoin from the blood compartment, but also simultaneously from the CSF, its toxic compartment.⁸³ This may be the reason why several reports describe a marked improvement in the patients' levels of consciousness during or following ECTR.

Despite the absence of robust evidence, the workgroup considered the following arguments in evaluating the risks and benefits of ECTR in phenytoin poisoning: the risk of prolonged coma with mechanical ventilation is not negligible; complications associated with ECTR are infrequent and usually mild; high efficiency intermittent ECTR can achieve rapid and substantial removal of phenytoin; and there is anecdotal evidence of clinical improvement following ECTR. Conversely, the mortality and long-term disability associated with phenytoin poisoning is very low; the cost of ECTR is not negligible, especially if the patient requires a transfer to another facility, and there is a theoretical risk of precipitating a seizure if phenytoin concentrations are abruptly lowered with ECTR in a patient with a known seizure disorder.

Given the lack of significant end-organ damage, the primary rationale for ECTR in phenytoin poisoning is to attenuate potential morbidity rather than to decrease related mortality. Several participants postulated that ECTR might decrease mechanical ventilation time, ICU length of stay and overall length of stay, which will in turn lessen financial cost; however, this effect would have to be weighed against the inherent risks and costs of ECTR mentioned above. Unfortunately, there are no cost-benefit studies that confirm or refute this hypothesis. Other participants believed that active supportive measures are sufficient. Taking into account the relative uncertainty concerning the toxicokinetic results detailed above, potential clinical benefit, risks, and the economic considerations and resource utilization, the workgroup proposed a neutral recommendation on the use of ECTR in severe phenytoin poisoning, meaning that ECTR would be reasonable in the right context. Twelve participants voted for ECTR, 8 participants supported a neutral position, and 7 voted against ECTR (median vote = 5, disagreement index < 1). Therefore, ECTR should probably only be considered in those patients that present following a massive ingestion who exhibit life-threatening toxicity and/or are expected to have very prolonged symptoms and in whom ECTR is considered to be safe. This case can be made for ECTR in a profoundly symptomatic patient, who by virtue of phenytoin's zero order kinetics, is likely to have a prolonged hospital stay. The opposite decision can be made for a moderately symptomatic patient needing a transfer to another center to receive ECTR, where supportive management can be preferred over ECTR. In other words, ECTR can be considered in selected cases where the potential benefit seems to outweigh the risks. Further study is needed to determine the place of ECTR in the management of phenytoin toxicity in the subpopulation with decreased protein binding of phenytoin (e.g., kidney failure, hypoalbuminemia).

2. Indications for ECTR

- **ECTR is suggested if prolonged coma is present or expected (2D)**
- **ECTR would be reasonable if prolonged incapacitating ataxia is present or expected (3D)**
- **We recommend NOT to perform ECTR solely based on a suspected dose of phenytoin ingested (1D)**
- **We recommend NOT to perform ECTR solely based on serum phenytoin concentration (1D)**

Rationale: The workgroup proposed that indications for ECTR initiation in any poisoning should be based on criteria that include exposure route (e.g., ingestion, intravenous), measurement of toxin in body fluids, technical examinations, and clinical symptoms and signs.

The workgroup agreed that there were too many uncertainties related to the dose of phenytoin ingested to initiate ECTR based on this information alone, certainly for a toxin that usually results in minimal or no long-term damage. The preferred management for patients presenting after an acute phenytoin exposure includes supportive measures, proper gastrointestinal decontamination with single dose activated charcoal and possibly MDAC. If the ingestion history is confirmed and the clinician suspects that major toxicity might ensue, then early communication with a toxicologist and nephrologist for consideration of possible ECTR may be warranted. This may be impossible in the event of IV overdose, given the rapid absorption and distribution (within minutes) and ensuing toxicity from the diluent.

Monitoring of serum phenytoin concentrations can confirm an acute exposure and may be available in a time frame short enough to guide clinical decisions. Nevertheless, the workgroup

suggested that the decision to initiate ECTR should be more dependent on symptomatology than on an arbitrary serum phenytoin concentration threshold. Prophylactic ECTR (i.e., ECTR before the appearance of symptoms) can be considered in poisons where irreversible or life-threatening clinical toxicity is expected (e.g., methanol, theophylline) but the workgroup did not endorse this approach for phenytoin.

Signs and symptoms following phenytoin poisoning are primarily neurologic. As stated earlier, phenytoin poisonings generally have a good prognosis and should be managed with supportive therapy. Coma in phenytoin poisoning is not due to a structural CNS lesion and it is also not considered life threatening in and of itself. However, coma following phenytoin poisoning may be prolonged and might necessitate protracted mechanical ventilation and ICU stays, warranting consideration of the risks of complications of prolonged intubation and/or intensive care treatment. For that reason, ECTR was not strongly recommended for coma by the workgroup. ECTR seems indicated only for an expected prolonged coma with the rationale to attenuate coma-induced complications and resource utilization.

There was less support for ECTR in prolonged incapacitating ataxia. These patients can easily be managed in a non-intensive care setting, and therefore would not use excessive resources, thus undermining the economic considerations. More benign symptoms like nystagmus did not warrant ECTR.

Seizures may occur in phenytoin poisoning and are very difficult to differentiate from seizures in a patient with seizure disorder. Some participants advocated the use of alternative anticonvulsants instead of ECTR in these patients, while other participants drew attention to the risk of dialyzing not only phenytoin, but also other anticonvulsants in patients with seizure disorders. No

agreement was reached on the use of ECTR in a patient with phenytoin poisoning if multiple seizures occur.

3. Cessation of ECTR

- **ECTR should be discontinued when clinical improvement is apparent (1D)**

Rationale: Because the recommendations for ECTR initiation are solely based on clinical symptoms, it is logical and reasonable to pursue ECTR until clear clinical improvement is present. Given the relatively modest clearances obtained with high-efficiency ECTRs, prolonged ECTR or a repeat session may be required. Phenytoin concentrations may rebound after ECTR, especially after a high-efficiency procedure.^{19,33,72,75,84} Although this is rarely a concern if caused by redistribution from deeper compartments, it may cause clinical morbidity if rebound is related to ongoing absorption, which has been reported as extensive in some cases.³³ It is therefore proposed to monitor clinical status and phenytoin concentrations serially over 24 hours after ECTR to help assess the need for subsequent sessions.

4. Choice of ECTR

- **Intermittent hemodialysis is the preferred ECTR in phenytoin poisoning (1D)**
- **Intermittent hemoperfusion is an acceptable alternative if intermittent hemodialysis is not available (1D)**

Rationale: According to the workgroup, intermittent HD is the preferred modality of ECTR in phenytoin poisoning. This recommendation is supported by the following arguments: clearance of

phenytoin has increased dramatically with the use of high-flux synthetic membranes compared to less efficient cuprophane or polyacrylonitrile filters;¹⁰³ intermittent HD is the most widely available modality of dialysis worldwide; more physicians and nurses are experienced with HD, with lesser risks of delay and uncertainty; the complication rate with HD appears favorable in comparison to HP, especially with regard to thrombocytopenia during HP, as described in some of the patients included in the cohort;^{18,73,88} the cost of HD favors it over HP. This is largely explained by the cost of monitoring and treating complications as well as the lower cost of dialysis filters versus charcoal cartridges, which need to be replaced regularly because of saturation of their adsorptive capacity.

Based on the fact that phenytoin is highly protein bound, a more efficient membrane and the optimization of both blood flow and effluent (dialysate and/or ultrafiltration) flow will have relatively minor but nevertheless significant effects on phenytoin clearance and are recommended.^{91,100,110,115}

If HD is not available, the workgroup recommended HP as an acceptable and useful alternative, as there is reliable data on its efficacy. HD and HP can also be used simultaneously in series with some clinical benefit.^{86,87} Peritoneal dialysis, albumin dialysis, exchange transfusion (ET) and therapeutic plasma exchange would not offer comparable results to HD or HP and are currently not recommended by EXTRIP for phenytoin poisoning. Continuous techniques, offer markedly lower clearances and removal rates compared to intermittent techniques.

CONCLUSION

The EXTRIP workgroup presents its recommendations for extracorporeal treatments in phenytoin poisoning. The great majority of cases can be treated with supportive care that may include single or multiple dose activated charcoal. In patients in whom prolonged coma is present

or expected, ECTR is suggested to accelerate elimination of phenytoin and to theoretically reduce ICU stay and its associated morbidity. The preferred choice of ECTR is high-efficiency intermittent hemodialysis. The workgroup advises to weigh the costs and risks associated with ECTR against the possible benefits in phenytoin toxicity and to individualize decisions to perform ECTR.

Acknowledgements:

The EXTRIP Workgroup also includes the following members: Ashish Bhalla, Diane P. Calello, Paul I. Dargan, Brian S. Decker, Tais F. Galvao, David S. Goldfarb, Lotte C. Hoegberg, David N. Juurlink, Jan T. Kielstein, Martin Laliberté, Yi Li, Kathleen D. Liu, Robert MacLaren, Robert Mactier, Bruno Mégarbane, Véronique Phan, Darren M. Roberts, Kevin M. Sowinski, Timothy J. Wiegand, James F. Winchester, and Christopher Yates.

We would like to acknowledge the tremendous work of our many colleagues who translated manuscripts and extracted data. All EXTRIP participants are listed at www.extrip-workgroup.org. We also acknowledge the important contribution from our librarians and other aids: Marc Lamarre, David Soteros, Salih Topal, Henry Gaston, and Brenda Gallant.

Table 1. Physicochemical and pharmacokinetic properties of phenytoin

Molecular mass	252 Da
Oral bioavailability	90%
Protein binding	90% (70-80% in hypoalbuminemia)
Volume of distribution	0.6-0.8 L/kg
Therapeutic range ^a	10-20 mcg/mL (39.6-79.2 µmol/L)
Toxic ingestion	≥ 20 mg/kg
Toxic plasma concentrations	≥ 20 mcg/mL (79 µmol/L)

^aTo convert units 1.0 mcg/mL = 3.96 µmol/L

Table 2. Strength of recommendation and level of evidence scaling for clinical outcomes

Strength of recommendation (consensus-based)	Level of evidence (based on GRADE system)
<p>Level 1 = Strong recommendation “We recommend...” <i>The course of action is considered appropriate by the large majority of experts with no major dissension The panel is confident that the desirable effects of adherence to the recommendation outweigh the undesirable effects.</i></p> <p>Level 2 = Weak recommendation “We suggest...” <i>The course of action is considered appropriate by the majority of experts but some degree of dissension exists amongst the panel. The desirable effects of adherence to the recommendation probably outweigh the undesirable effects.</i></p> <p>Level 3 = Neutral recommendation “It would be reasonable...” <i>The course of action could be considered appropriate in the right context</i></p> <p>No recommendation <i>No agreement was reached by the group of experts</i></p>	<p>Grade A = High level of evidence <i>The true effect lies close to our estimate of the effect</i></p> <p>Grade B = Moderate level of evidence <i>The true effect is likely to be close to our estimate of the effect, but there is a possibility that it is substantially different</i></p> <p>Grade C = Low level of evidence <i>The true effect may be substantially different from our estimate of the effect</i></p> <p>Grade D = Very low level of evidence <i>Our estimate of the effect is just a guess, and it is very likely that the true effect is substantially different from our estimate of the effect</i></p>

Adapted from Lavergne V, Nolin TD, Hoffman RS, et al. The EXTRIP (EXtracorporeal TReatments In Poisoning) workgroup: guideline methodology. *Clin Toxicol (Phila)*. 2012;50(5):403-413.

Table 3. Aggregate clinical outcomes of the 31 overdose patients described in case reports or case series

Demographics	Age in years [mean (range)]	26.7 (0.1-77)
	Male (%)	63.3%
Phenytoin exposure	Amount ingested (grams)	6.8 (1-21.5)
	Peak total phenytoin conc [mean (range)] (mcg/mL)	69.8 (15-200.7)
	Time from ingestion to presentation (hours)	18.1 (1-120)
Toxic symptoms (%)	Altered consciousness	90.3%
	Seizure (one or more)	19.4%
	Respiratory failure	32.3%
	Dysarthria/ataxia	25.8%
Other treatments (%)	Mechanical ventilation	25.8%
	MDAC	9.7%
ECTR (N)	HD	7
	HP	7
	CRRT	0
	HP-HD	3
	TPE	3
	ET	1
	PD	6
	Liver support therapy (MARS)	1
	More than 1 ECTR	3
Outcome (%)	Sequelae	12.9%
	Death	6.5%

HD, intermittent hemodialysis; HP, hemoperfusion; CRRT, continuous renal replacement therapy; HD-HP, intermittent hemodialysis and hemoperfusion in series; TPE, therapeutic plasma exchange; ET, exchange transfusion; PD, peritoneal dialysis; MARS, molecular adsorbent recirculating system.

Table 4. Clearance of various ECTRs from accepted articles

ECTR	Patients	Clearance (mL/min)		References
		Average	Range	
Conventional HD	10	16.7	6.5-42	71,90-92
High efficiency HD	3	68.1	44.3-112	19,86,103
Charcoal HP	4	28.8	18-42	72,73,83,86
HD-HP	1	58.4		86
TPE	14	18.5	7.8-43	75,78,80,81,93,95-99,102
PD	6	3.3	0.2-10.6	36,68,69,74,76,94
ET	1	3.1		76
CRRT	4	6.5	0.9-13	100,104

HD, intermittent hemodialysis; HP, hemoperfusion; HD-HP, intermittent hemodialysis and hemoperfusion in series; TPE, therapeutic plasma exchange; PD, peritoneal dialysis; ET, exchange transfusion; CRRT, continuous renal replacement therapies.

Table 5. Toxicokinetic grading attributed to individual patients

	Number of Patients Graded								
	PD	HP	HD (conv)	HD (high-eff)	HD-HP (in series)	CRRT	TPE	ET	LST
Dialyzable		2		2	1				1
Moderately dialyzable		2		2			7		
Slightly dialyzable	1		5			2	7	1	
Not dialyzable	5		4			2	2		

PD, peritoneal dialysis; HP, hemoperfusion; HD (conv), conventional intermittent hemodialysis; HD (high-eff), high efficiency intermittent hemodialysis; HD-HP, intermittent hemodialysis and hemoperfusion in series; CRRT, continuous renal replacement therapy; TPE, therapeutic plasma exchange; ET, exchange transfusion; LST, liver support therapy.

Box 1. Executive summary of recommendations

General statement regarding use of ECTR

- ECTR would be reasonable in selected cases of severe phenytoin poisoning (3D)

Indications for ECTR

- ECTR is suggested if prolonged coma is present or expected (2D)
- ECTR would be reasonable if prolonged incapacitating ataxia is present or expected (3D)
- We recommend NOT to perform ECTR solely based on suspected dose of phenytoin ingested (1D)
- We recommend NOT to perform ECTR solely based on serum phenytoin concentration (1D)

Cessation of ECTR

- ECTR should be discontinued when clinical improvement is apparent (1D)


Choice of ECTR

- Intermittent HD is the preferred ECTR in phenytoin poisoning (1D)
- Intermittent HP is an acceptable alternative if intermittent HD is not available (1D)

Figure legend

Figure 1. Summary of literature search on use of ECTR in phenytoin poisoning.

Figure 1.



Supplementary Material

Item S1. Represented Societies

Item S1. Societies represented in EXTRIP

American Academy of Clinical Toxicology
American College of Emergency Physicians
American College of Medical Toxicology
American Society of Nephrology
American Society of Pediatric Nephrology
Asia Pacific Association of Medical Toxicology
Australian and New Zealand Intensive Care Society
Australian and New Zealand Society of Nephrology
Brazilian Association of Poison Control Centers and Clinical Toxicologists
Brazilian Society of Nephrology
Brazilian Society of Toxicology
Canadian Association of Poison Control Centres
Canadian Association of Emergency Physicians
Canadian Society of Nephrology
Chinese College of Emergency Physicians
Chinese Medical Doctor Association
European Association of Poison Centres and Clinical Toxicologists
European Renal Best Practice
European Society of Emergency Medicine
European Society of Intensive Care Medicine
French Language Society of Resuscitation
German Society of Nephrology
International Pediatric Nephrology Association
International Society of Nephrology
Latin American Society of Nephrology and Hypertension
National Kidney Foundation
Pediatric Continuous Renal Replacement Therapy
Pediatric Critical Care Medicine
Quebec Association of Emergency Physicians
Quebec Association of Specialists in Emergency Medicine
Quebec Society of Nephrology
Renal Association
Society of Critical Care Medicine
Spanish Clinical Toxicology Foundation

References

1. Ghannoum M, Nolin TD, Lavergne V, Hoffman RS. Blood purification in toxicology: nephrology's ugly duckling. *Adv Chronic Kidney Dis*. 2011;18(3):160-166.
2. Lavergne V, Nolin TD, Hoffman RS, et al. The EXTRIP (EXtracorporeal TReatments In Poisoning) workgroup: guideline methodology. *Clin Toxicol (Phila)*. 2012;50(5):403-413.
3. Ghannoum M, Nolin TD, Goldfarb DS, et al. Extracorporeal Treatment for Thallium Poisoning: Recommendations from the EXTRIP Workgroup. *Clin J Am Soc Nephrol*. 2012;7(10):1682-1690.
4. Yates C, Galvao T, Sowinski KM, et al. Extracorporeal treatment for tricyclic antidepressant poisoning: recommendations from the EXTRIP Workgroup. *Semin Dial*. 2014;27(4):381-389.
5. Mactier R, Laliberte M, Mardini J, et al. Extracorporeal treatment for barbiturate poisoning: recommendations from the EXTRIP Workgroup. *Am J Kidney Dis*. 2014;64(3):347-358.
6. Roberts DM, Yates C, Megarbane B, et al. Recommendations for the role of extracorporeal treatments in the management of acute methanol poisoning: a systematic review and consensus statement. *Crit Care Med*. 2015;43(2):461-472.
7. Gosselin S, Juurlink DN, Kielstein JT, et al. Extracorporeal treatment for acetaminophen poisoning: recommendations from the EXTRIP workgroup. *Clin Toxicol (Phila)*. 2014;52(8):856-867.

8. Ghannoum M, Yates C, Galvao TF, et al. Extracorporeal treatment for carbamazepine poisoning: Systematic review and recommendations from the EXTRIP workgroup. *Clin Toxicol (Phila)*. 2014;52(10):993-1004.
9. Lavergne V, Ouellet G, Bouchard J, et al. Guidelines for reporting case studies on extracorporeal treatments in poisonings: methodology. *Semin Dial*. 2014;27(4):407-414.
10. Ghannoum M, Wiegand TJ, Liu KD, et al. Extracorporeal treatment for theophylline poisoning: Systematic review and recommendations from the EXTRIP workgroup. *Clin Toxicol (Phila)*. 2015;53(4):215-229.
11. Decker BS, Goldfarb DS, Dargan PI, et al. Extracorporeal Treatment for Lithium Poisoning: Systematic Review and Recommendations from the EXTRIP Workgroup. *Clin J Am Soc Nephrol*. 2015;10(5):875-887.
12. Calello DP, Liu KD, Wiegand TJ, et al. Extracorporeal Treatment for Metformin Poisoning: Systematic Review and Recommendations From the Extracorporeal Treatments in Poisoning Workgroup. *Crit Care Med*. 2015;43(8):1716-1730.
13. Ghannoum M, Laliberte M, Nolin TD, et al. Extracorporeal treatment for valproic acid poisoning: Systematic review and recommendations from the EXTRIP workgroup. *Clin Toxicol (Phila)*. 2015;53(5):454-465.
14. Woodbury DM, Kemp JW. Pharmacology and mechanisms of action of diphenylhydantoin. *Psychiatr Neurol Neurochir*. 1971;74(2):91-115.
15. Rand BO, Kelly WA, Ward AA, Jr. Electrophysiological studies of the action of intravenous diphenylhydantoin (Dilantin). *Neurology*. 1966;16(10):1022-1032.
16. Dalessio DJ. Medical treatment of the major neuralgias. *Semin Neurol*. 1988;8(4):286-290.

17. Pincus JH, Grove I, Marino BB, Glaser GE. Studies on the mechanism of action of diphenylhydantoin. *Arch Neurol.* 1970;22(6):566-571.
18. Kawasaki C, Nishi R, Uekihara S, Hayano S, Otagiri M. Charcoal hemoperfusion in the treatment of phenytoin overdose. *Am J Kidney Dis.* 2000;35(2):323-326.
19. Ghannoum M, Troyanov S, Ayoub P, Lavergne V, Hewlett T. Successful hemodialysis in a phenytoin overdose: case report and review of the literature. *Clin Nephrol.* 2010;74(1):59-64.
20. Ramasamy K, Narayan SK, Shewade DG, Chandrasekaran A. Influence of CYP2C9 genetic polymorphism and undernourishment on plasma-free phenytoin concentrations in epileptic patients. *Ther Drug Monit.* 2010;32(6):762-766.
21. Richens A. Clinical pharmacokinetics of phenytoin. *Clin Pharmacokinet.* 1979;4(3):153-169.
22. van Peer AP, Belpaire FM, Bogaert MG. Distribution of phenytoin and phenylbutazone in normal and uraemic rabbits. *Arch Int Pharmacodyn Ther.* 1978;234(2):348.
23. Masur H, Elger CE, Ludolph AC, Galanski M. Cerebellar atrophy following acute intoxication with phenytoin. *Neurology.* 1989;39(3):432-433.
24. Kuruvilla T, Bharucha NE. Cerebellar atrophy after acute phenytoin intoxication. *Epilepsia.* 1997;38(4):500-502.
25. Chaikin P, Adir J. Unusual absorption profile of phenytoin in a massive overdose case. *J Clin Pharmacol.* 1987;27(1):70-73.
26. Rosenberg J, Benowitz NL, Pond S. Pharmacokinetics of drug overdose. *Clin Pharmacokinet.* 1981;6(3):161-192.

27. Harrison DC, Meffin PJ, Winkle RA. Clinical pharmacokinetics of antiarrhythmic drugs. *Prog Cardiovasc Dis.* 1977;20(3):217-242.
28. Brandolese R, Scordo MG, Spina E, Gusella M, Padriani R. Severe phenytoin intoxication in a subject homozygous for CYP2C9*3. *Clin Pharmacol Ther.* 2001;70(4):391-394.
29. Mowry JB, Spyker DA, Cantilena LR, Jr., McMillan N, Ford M. 2013 Annual Report of the American Association of Poison Control Centers' National Poison Data System (NPDS): 31st Annual Report. *Clin Toxicol (Phila).* 2014;52(10):1032-1283.
30. Kutt H, Winters W, Kokenge R, McDowell F. Diphenylhydantoin Metabolism, Blood Levels, and Toxicity. *Arch Neurol.* 1964;11:642-648.
31. Raines A, Niner JM, Pace DG. A comparison of the anticonvulsant, neurotoxic and lethal effects of diphenylbarbituric acid, phenobarbital and diphenylhydantoin in the mouse. *J Pharmacol Exp Ther.* 1973;186(2):315-322.
32. Case log counts (Generic). Enterprise Reports, National Poison Data System, American Association of Poison Control Centers. <http://www.aapcc.org/data-system/>. Accessed May 15, 2015.
33. Miller MA, Crystal CS, Patel MM. Hemodialysis and hemoperfusion in a patient with an isolated phenytoin overdose. *Am J Emerg Med.* 2006;24(6):748-749.
34. Mowry JB, Spyker DA, Cantilena LR, Jr., Bailey JE, Ford M. 2012 annual report of the American association of poison control centers' national poison data system (NPDS): 30th annual report. *Clin Toxicol (Phila).* 2013;51(10):949-1229.
35. Gerstman J, King A, Menke N, Johnston J, Lynch M, Pizon AF. Hemodialysis as a treatment for severe phenytoin toxicity. *Clin Toxicol (Phila).* 2013;51(7):575-724.

36. Narcy P, Zorza G, Taburet AM, Mersch JM, Devictor D, Huault G. Severe poisoning with intravenous phenytoin in the newborn. Value of peritoneal dialysis [in French]. *Arch Fr Pediatr*. 1990;47(8):591-593.
37. Sen S, Ratnaraj N, Davies NA, et al. Treatment of phenytoin toxicity by the molecular adsorbents recirculating system (MARS). *Epilepsia*. 2003;44(2):265-267.
38. Leuschner F, Neumann W, Reith H. A contribution to pharmacology and toxicology of injectable phenytoin solutions [in German]. *Arzneimittelforschung*. 1977;27(4):811-819.
39. Pillai U, Hothi J, Bhat Z. Severe propylene glycol toxicity secondary to use of anti-epileptics. *Am J Ther*. 2014;21(4):e106-e109.
40. Watson WA, Litovitz TL, Rodgers GC, Jr., et al. 2002 annual report of the American Association of Poison Control Centers Toxic Exposure Surveillance System. *Am J Emerg Med*. 2003;21(5):353-421.
41. Presutti M, Pollet L, Stordeur JM, Bruder N, Gouin F. Acute poisoning with phenytoin caused by an error in the administration of fosphenytoin (Prodilantin) [in French]. *Ann Fr Anesth Reanim*. 2000;19(9):688-690.
42. McBryde KD, Wilcox J, Kher KK. Hyperphosphatemia due to fosphenytoin in a pediatric ESRD patient. *Pediatr Nephrol*. 2005;20(8):1182-1185.
43. Keegan MT, Bondy LR, Blackshear JL, Lanier WL. Hypocalcemia-like electrocardiographic changes after administration of intravenous fosphenytoin. *Mayo Clin Proc*. 2002;77(6):584-586.
44. Grageda M, Saini AP, Trout LC, Cyran SE, Halstead ES. Severe cardiomyopathy in an infant after iatrogenic fosphenytoin overdose. *Clin Pediatr (Phila)*. 2014;53(8):791-793.

45. Adams BD, Buckley NH, Kim JY, Tipps LB. Fosphenytoin may cause hemodynamically unstable bradydysrhythmias. *J Emerg Med.* 2006;30(1):75-79.
46. Curtis DL, Piibe R, Ellenhorn MJ, Wasserberger J, Ordog G. Phenytoin toxicity: predictors of clinical course. *Vet Hum Toxicol.* 1989;31(2):162-163.
47. Curtis DL, Piibe R, Ellenhorn MJ, Wasserberger J, Ordog G. Phenytoin toxicity: a review of 94 cases. *Vet Hum Toxicol.* 1989;31(2):164-165.
48. Craig S. Phenytoin poisoning. *Neurocrit Care.* 2005;3(2):161-170.
49. Chyka PA, Seger D, Krenzelok EP, et al. Position paper: single-dose activated charcoal. *Clin Toxicol (Phila).* 2005;43(2):61-87.
50. Skinner CG, Chang AS, Matthews AS, Reedy SJ, Morgan BW. Randomized controlled study on the use of multiple-dose activated charcoal in patients with supratherapeutic phenytoin levels. *Clin Toxicol (Phila).* 2012;50(8):764-769.
51. Mauro LS, Mauro VF, Brown DL, Somani P. Enhancement of phenytoin elimination by multiple-dose activated charcoal. *Ann Emerg Med.* 1987;16(10):1132-1135.
52. Rowden AM, Spoor JE, Bertino JS, Jr. The effect of activated charcoal on phenytoin pharmacokinetics. *Ann Emerg Med.* 1990;19(10):1144-1147.
53. Mofenson HC, Caraccio TR, Greensher J, D'Agostino R, Rossi A. Gastrointestinal dialysis with activated charcoal and cathartic in the treatment of adolescent intoxications. *Clin Pediatr (Phila).* 1985;24(12):678-684.
54. Pettrey J, Rose R, Wills B, Cumpston K. Activated charcoal does not reduce duration of phenytoin toxicity in hospitalized patients [abstract]. *Clin Toxicol (Phila).* 2011;49:536.

55. Rivera JV, Aaronson P, Schauben J, Sollee D, Kunisaki T. Clinical outcomes after utilization of multi-dose activated charcoal in phenytoin toxicity [abstract]. *Clin Toxicol (Phila)*. 2014;52(7):752.
56. Dolgin JG, Nix DE, Sanchez J, Watson WA. Pharmacokinetic simulation of the effect of multiple-dose activated charcoal in phenytoin poisoning--report of two pediatric cases. *Drug Intell Clin Pharm*. 1991;25(6):646-649.
57. Vale J, Krenzelok, EP., Barceloux VD. Position statement and practice guidelines on the use of multi-dose activated charcoal in the treatment of acute poisoning. American Academy of Clinical Toxicology; European Association of Poisons Centres and Clinical Toxicologists. *J Toxicol Clin Toxicol*. 1999;37(6):731-751.
58. Su M, Amitai A. Phenytoin poisoning. In: Basow D, ed. UpToDate. Waltham, MA; 2014. http://www.uptodate.com/contents/phenytoin-poisoning?source=search_result&search=phenytoin+poisoning&selectedTitle=1%7E25. Accessed March 15th, 2015.
59. Miller C, Joyce DM. Phenytoin toxicity. In: WebMD, ed. Medscape. New York; 2012. <http://emedicine.medscape.com/article/816447-treatment#a1126>. Accessed March 15th, 2015.
60. Fountain J. Phenytoin. In: Centre NP, ed. ToxinZ. New Zealand; 2014. <http://toxinZ.com/>. Accessed March 15th, 2015.
61. Doyon S. Anticonvulsants. In: Nelson LS, Lewin NA, Howland MA, Hoffman RS, Goldfrank LR, Flomenbaum N, eds. *Goldfrank's Toxicologic Emergencies*. New York, NY: McGraw-Hill; 2011:698-710.

62. Phenytoin. Wikitox. <http://curriculum.toxicology.wikispaces.net/2.1.11.2.1+Phenytoin>. Accessed March 15th, 2015.
63. Appraisal of guidelines for research & evaluation. AGREE instrument. In: School SGsHM, ed. London: The AGREE Collaboration; 2001.
64. Atkins D, Best D, Briss PA, et al. Grading quality of evidence and strength of recommendations. *BMJ*. 2004;328(7454):1490.
65. Fitch K, Bernstein SJ, Aguilar MD, et al. The RAND/UCLA Appropriateness Method User's Manual. In: RAND, ed. Santa Monica, USA: RAND; 2011:
<http://www.rand.org/>.
66. Schreiner GE. The role of hemodialysis (artificial kidney) in acute poisoning. *AMA Arch Intern Med*. 1958;102(6):896-913.
67. Theil G, Richter R, Powell M, Doolaan P. Acute dilantin poisoning. *Neurology*. 1961;11:138-142.
68. Andia J, Westphal M, Anthone R, Anthone S. Severe, acute diphenylhydantoin intoxication treated with peritoneal lavage. *N Y State J Med*. 1968;68(13):1861-1863.
69. Blair AA, Hallpike JF, Lascelles PT, Wingate DL. Acute diphenylhydantoin and primidone poisoning treated by peritoneal dialysis. *J Neurol Neurosurg Psychiatry*. 1968;31(5):520-523.
70. Tenckhoff H, Sherrard DJ, Hickman RO, Ladda RL. Acute diphenylhydantoin intoxication. *Am J Dis Child*. 1968;116(4):422-425.
71. Rubinger D, Levy M, Roll D, Czaczkes JW. Inefficiency of haemodialysis in acute phenytoin intoxication. *Br J Clin Pharmacol*. 1979;7(4):405-407.

72. Baehler RW, Work J, Smith W, Dominic JA. Charcoal hemoperfusion in the therapy for methsuximide and phenytoin overdose. *Arch Intern Med.* 1980;140(11):1466-1468.
73. Crome P, Hampel G, Widdop B, Goulding R. Experience with cellulose acetate-coated activated charcoal haemoperfusion in the treatment of severe hypnotic drug intoxication. *Postgrad Med J.* 1980;56(661):763-766.
74. Czajka PA, Anderson WH, Christoph RA, Banner W, Jr. A pharmacokinetic evaluation of peritoneal dialysis for phenytoin intoxication. *J Clin Pharmacol.* 1980;20(10):565-569.
75. Bambauer R, Jutzler GA, Stolz D, Doenecke P. Detoxication using a technically simplified plasma filtration system [in German]. *Intensivmedizin.* 1982;19 (2):79-82.
76. Lindahl S, Westerling D. Detoxification with peritoneal dialysis and blood exchange after diphenylhydantoin intoxication. *Acta Paediatr Scand.* 1982;71(4):665-666.
77. Durakovic Z, Plavsic F. Use of hemodialysis in the treatment of phenobarbitone and phenytoin poisoning [in Croatian]. *Arh Hig Rada Toksikol.* 1986;37(1):59-66.
78. Larsen LS, Sterrett JR, Whitehead B, Marcus SM. Adjunctive therapy of phenytoin overdose--a case report using plasmapheresis. *J Toxicol Clin Toxicol.* 1986;24(1):37-49.
79. Springer W, Greiner C, Wehran HJ. Successful hemodialysis/hemoperfusion in a small child intoxicated by a mixture of phenytoin and carbamezepine [in German]. *Z Klin Med.* 1987;42(16):1401-1402.
80. Ogura M, Ohbu S, Ohiwa T, Hayashida N, Honda M. A successful case of hollow-fiber plasmapheresis for prolonged phenytoin intoxication [in Japanese]. *Nippon Naika Gakkai Zasshi.* 1989;78(8):1194-1198.
81. Scharf J, Ruder H, Harms D. Plasma separation in acute, intravenous phenytoin poisoning [in German]. *Monatsschr Kinderheilkd.* 1990;138(4):227-230.

82. Morikawa N, Mori K, Fujii I, Takeyama M. Pharmacokinetic study in acute antiepileptic intoxication treated with hemodialysis and hemoperfusion [in Japanese]. *Japan J Clin Pharmacol Ther.* 1992;23(2):469-474.
83. Kanayama Y, Itakura Y, Iwasaki M, et al. Changes in phenytoin concentrations in blood and cerebrospinal fluid caused by direct hemoperfusion in a patient intoxicated with phenytoin. *Ther Apher.* 1998;2(1):74-77.
84. Craig S. Phenytoin overdose complicated by prolonged intoxication and residual neurological deficits. *Emerg Med Australas.* 2004;16(4):361-365.
85. Sung SF, Chiang PC, Tung HH, Ong CT. Charcoal hemoperfusion in an elderly man with life-threatening adverse reactions due to poor metabolism of phenytoin. *J Formos Med Assoc.* 2004;103(8):648-652.
86. De Schoenmakere G, De Waele J, Terryn W, et al. Phenytoin intoxication in critically ill patients. *Am J Kidney Dis.* 2005;45(1):189-192.
87. Eyer F, Felgenhauer N, Pfab R, Thurmel K, Zilker T. Treatment of severe intravenous phenytoin overdose with hemodialysis and hemoperfusion. *Med Sci Monit.* 2008;14(12):CS145-148.
88. Kumar PP, Lingappa L, Shah MA, Shaikh FA. Charcoal hemoperfusion for phenytoin intoxication. *Indian Pediatr.* 2012;49(2):152-153.
89. Swartzenburger GS, Raja AH, Lynch MJ, Menke NB. High-flux hemodialysis enhances phenytoin elimination and improves neurological function after oral overdose: A case report [abstract]. *Clin Toxicol (Phila).* 2014;52(7):763-764.
90. Adler DS, Martin E, Gambertoglio JG, Tozer TN, Spire JP. Hemodialysis of phenytoin in a uremic patient. *Clin Pharmacol Ther.* 1975;18(1):65-69.

91. Bosl R, Shideman JR, Meyer RM, Buselmeier TJ, von Hartitzsch B, Kjellstrand CM. Effects and complications of high efficiency dialysis. *Nephron*. 1975;15(2):151-160.
92. Martin E, Gambertoglio JG, Adler DS, Tozer TN, Roman LA, Grausz H. Removal of phenytoin by hemodialysis in uremic patients. *JAMA*. 1977;238(16):1750-1753.
93. Liu E, Rubenstein M. Phenytoin removal by plasmapheresis in thrombotic thrombocytopenic purpura. *Clin Pharmacol Ther*. 1982;31(6):762-765.
94. Hays DP, Primack WA, Abroms IF. Phenytoin clearance by continuous ambulatory peritoneal dialysis. *Drug Intell Clin Pharm*. 1985;19(6):429-431.
95. Nasca TJ, Huff N, Livengood BH. Phenytoin pharmacokinetics in therapeutic plasmapheresis. *J Clin Pharmacol*. 1985;25(4):302-304.
96. Brown-Molnar CS, Bauer LA, Horn JR. Phenytoin removal by plasmapheresis in renal insufficiency. *Am J Nephrol*. 1986;6(4):302-306.
97. Silberstein LE, Shaw LM. Effect of plasma exchange on phenytoin plasma concentration. *Ther Drug Monit*. 1986;8(2):172-176.
98. White RL, Garnett WR, Allen JH. Phenytoin removal during plasma exchange. *J Clin Apher*. 1987;3(3):147-150.
99. Tobias JD, Baker DK, Hurwitz CA. Removal of phenytoin by plasmapheresis in a patient with thrombotic thrombocytopenia purpura. *Clin Pediatr (Phila)*. 1992;31(2):105-108.
100. Lau AH, Kronfol NO. Effect of continuous hemofiltration on phenytoin elimination. *Ther Drug Monit*. 1994;16(1):53-57.
101. Olsen KM, Marx MA, Monaghan MS, Barnes E, Ackerman BH, Pappas AA. Phenytoin and plasmapheresis: importance of sampling times and impact of obesity. *Ther Drug Monit*. 1994;16(6):624-628.

- 102.** Mansur LI, Murrow RW, Garrelts JC, Minns GO. Rebound of plasma free phenytoin concentration following plasmapheresis in a patient with thrombotic thrombocytopenic purpura. *Ann Pharmacother.* 1995;29(6):592-595.
- 103.** Frenchie D, Bastani B. Significant removal of phenytoin during high flux dialysis with cellulose triacetate dialyzer. *Nephrol Dial Transplant.* 1998;13(3):817-818.
- 104.** Oltrogge KM, Peppard WJ, Saleh M, Regner KR, Herrmann DJ. Phenytoin removal by continuous venovenous hemofiltration. *Ann Pharmacother.* 2013;47(9):1218-1222.
- 105.** Bezzaoucha S, Merghoub A, Lamarche C, et al. Hemodialysis effects on phenytoin pharmacokinetics. *Eur J Clin Pharmacol.* 2014;70(4):499-500.
- 106.** Aweeka FT, Gottwald MD, Gambertoglio JG, et al. Pharmacokinetics of fosphenytoin in patients with hepatic or renal disease. *Epilepsia.* 1999;40(6):777-782.
- 107.** Klehr HU, Bley T, Platzbecker S, Schwickart R. Influence of charcoal hemoperfusion on the elimination of diphenylhydantoin, phenobarbital and isoniazid in experimental drug poisoning [abstract]. *Artif Organs.* 1979;3 (3):303.
- 108.** Kroh UF, Dinges GK, Lukasewitz P, Steinhausser WU, Holl TJ, Lennartz H. Elimination of drugs by the new polyamide hemofilter FH77H during various in vitro conditions. *Blood Purif.* 1998;16(1):49-56.
- 109.** Reiter K, Bordonni V, Dall'Olio G, et al. In vitro removal of therapeutic drugs with a novel adsorbent system. *Blood Purif.* 2002;20(4):380-388.
- 110.** Churchwell MD, Pasko DA, Smoyer WE, Mueller BA. Enhanced clearance of highly protein-bound drugs by albumin-supplemented dialysate during modeled continuous hemodialysis. *Nephrol Dial Transplant.* 2009;24(1):231-238.

- 111.** Mellick LB, Morgan JA, Mellick GA. Presentations of acute phenytoin overdose. *Am J Emerg Med.* 1989;7(1):61-67.
- 112.** Nauth-Misir TN. A case of gross overdosage of soluble phenytoin. *Br Med J.* 1948;2(4578):646.
- 113.** Ouellet G, Bouchard J, Ghannoum M, Decker BS. Available extracorporeal treatments for poisoning: overview and limitations. *Semin Dial.* 2014;27(4):342-349.
- 114.** Ghannoum M, Bouchard J, Nolin TD, Ouellet G, Roberts DM. Hemoperfusion for the treatment of poisoning: technology, determinants of poison clearance, and application in clinical practice. *Semin Dial.* 2014;27(4):350-361.
- 115.** Bouchard J, Roberts DM, Roy L, et al. Principles and operational parameters to optimize poison removal with extracorporeal treatments. *Semin Dial.* 2014;27(4):371-380.